

DATE _____

TRANSFER HOSP _____

PATIENT NAME _____

DESTINATION HOSP _____

AGE _____ DOB _____ SEX _____

EMS AGENCY _____

MEDICAL RECORDS # _____

EMS RUN # _____

ACCOUNT# _____

REGIONAL NEUROLOGICAL ABBREVIATED ASSESSMENT TOOL FOR TRANSFER

LKWT:		ABBREVIATED NIHSS*																
		TIME	HEART RATE	HEART RHYTHM	BLOOD PRESSURE SBP<180 and DBP<105	MOTOR SCORE				LEVEL OF CONSCIOUSNESS	DYSARTHRIA	LANGUAGE	TOTAL SCORE	ANGIO-EDEMA Y / N	INITIALS			
RIGHT ARM	LEFT ARM					RIGHT LEG	LEFT LEG											
Administration	VS Prior TPA																	
	tPA 15min																	
	tPA 15min																	
	tPA 15min																	
	tPA 15min																	
	POST tPA Q15-1HR	15min																
		15min																
		15min																
		15min																
		Q30 X 6 HOURS	30min															
			30min															
			30min															
			30min															
			30min															
			30min															
30min																		
30min																		
30min																		
30min																		
30min																		
30min																		
30min																		
30min																		
MOTOR SCORE 0 = No drift, arms hold 90 degrees (sitting) or 45 degrees (supine) for full 10 seconds; or leg holds 30 degrees for full 5 seconds. 1 = Drift, arm holds 90 degrees (sitting) or 45 degrees (supine), but drifts down before full 10 seconds; or leg holds 30 degrees but drifts down before 5 full seconds, but does not hit bed or other support. 2 = Some effort against gravity; arm cannot get to or maintain (if cued) 90 degrees (sitting) or 45 degrees (supine); or leg cannot get to or maintain 30 degrees, drifts down to bed, but has some effort against gravity. 3 = No effort against gravity, limb falls. 4 = No movement. 5 = Amputation; joint fusion.										LEVEL OF CONSCIOUSNESS 0 = Alert 1 = Not alert, arouses with minor stimulation 2 = Not alert, arouses with strong, repeated stimulation 3 = Responds with reflexes or is unresponsive			DYSARTHRIA 0 = Normal 1 = Mild to moderate; slurs but can be understood 2 = Severe; so slurred it is unintelligible 3 = Intubated or other physical barrier					
NOTE: 4-point increase from baseline should trigger further neurological assessment & notify neurologist STAT. Angio edema: Look for signs of unilateral or bilateral tongue enlargement q15min X 2hrs post (Alteplase).										BEST LANGUAGE 0 = No aphasia 1 = Mild to moderate; some loss but makes conversation with materials 2 = Severe aphasia; information exchanged is limited 3 = Mute; no usable speech or auditory comprehension								
STOP the infusion immediately, and notify neurologist or ED physician if the patient displays any of the following symptoms: headache, severe chest pain, profuse bleeding, flank pain, edema of mouth/ lips/ tongue, or seizures.																		
BAR CODE					COMMENTS (i.e., glaze, facial palsy, ataxia, sensory, changes in vital signs):													
					_____ _____ _____													

