

Geriatric Trauma References

Brar, SS, et al. Sodium Bicarbonate vs Sodium Chloride for the prevention of contrast medium-induced nephropathy in patients undergoing coronary angiography. *JAMA* 2008; 300(9):1038-1046. <http://jama.ama-assn.org/cgi/content/full/300/9/1038> Accessed November 12, 2012.

Brasel, KB, et al. Rib fractures: Relationship with pneumonia and mortality. *Crit Care Med.* 2006; 34(6): 1642-1646.

Brown, CVR, et al. Recombinant Factor VIIa for the Correction of Coagulopathy Before Emergent Craniotomy in Blunt Trauma Patients. *J Trauma.*2010; 68:348-352.

Bulger EM, et al. Rib fractures in the elderly. *J Trauma.*2000; 28:1040-1047.

Carrier, FM, et al. Effect of epidural analgesia in patients with traumatic rib fractures: a systematic review and meta-analysis of randomized controlled trials. *Can J Anaesth.*2009; 56(3):230-242.

Caterino JM, et al. Identification of an age cutoff for increased mortality in patients with elderly trauma. *Am J Emerg Med.*2010; 28:151–158.

Como, JJ, et al. Practice management guidelines for identification of cervical spine injuries following trauma: update from the Eastern Association for the Surgery of Trauma Practice Management Guidelines Committee. *J Trauma.*2009; 67:651-659.

Golob, JF, et al. Isolated cervical spine fractures in the elderly: a deadly injury. *J Trauma.*2008; 64:311-315.


Ivascu, FA, et al. Rapid warfarin reversal in anticoagulated patients with traumatic intracranial hemorrhage reduces hemorrhage progression and mortality. *J Trauma.* 2005, 59:1131–1139.

Kochanek, KD, et al. Deaths, preliminary data for 2009. *National Vital Statistics Reports.*2011; 59(4). http://www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_04.pdf Accessed November 12, 2012.

Mahoney, LK. Rib Fractures. Medscape Reference; updated June 2011. <http://emedicine.medscape.com/article/825981-overview>. Accessed November 12, 2012.

Majumdar, SR, et al. Forced euvolemic diuresis with mannitol and furosemide for prevention of contrast-induced nephropathy in patients with CKD undergoing coronary angiography: a randomized controlled trial. *Am J Kidney Dis* 2009; 54(4):602-9. Epub 2009 Jun 17 <http://www.ncbi.nlm.nih.gov/pubmed/19535188>. Accessed November 12, 2012.

Manion, SC & Brennan, TJ. Thoracic Epidural Analgesia and Acute Pain Management. *Anesthesiology.*2011; 115(1):181-188.



Schoenfeld, BW, et al. Type II odontoid fractures of the cervical spine. Do treatment type and medical comorbidities affect mortality in elderly patients? *Spine*.2011; 36(11):879-885.

Shifflette VK, et al. Should age be a factor to change from a level II to a level I trauma activation? *J Trauma*. 2010; 69:88 –92.

Tashjian, RZ, et al. Halo-vest immobilization increased early morbidity and mortality in elderly odontoid fractures. *J Trauma*. 2006; 60(1):199–203.

Zagler, A, et al. N-acetylcysteine and contrast-induced nephropathy: a meta-analysis of 13 randomized trials. *Am Heart J* 2006; 151(1):140-5.