

Pedi  $\leq 17$

Adult  $\geq 18, < 65$

Geri  $\geq 65$

**If any Red Criteria met, transport to Level 1 Trauma Center**

- R1** Patient not awake and appropriate
- R2** Active airway assistance required (ie. more than supplemental O2), or respiratory distress
- R3** Weak carotid/femoral pulse or absent distal pulses
- R4** BP <70 plus 2X Age (BP <90 age >10)
- R5** Pelvic instability or Chest wall instability or crepitus
- R6** Acute paralysis, loss of sensation, or suspected spinal cord injury
- R7** Amputation proximal to wrist or ankle
- R8**  $\geq 5\%$  BSA partial/full thickness burns
- R9** Penetrating injury to head (or depressed skull fracture), neck, torso, extremities proximal to elbow or knee, excluding superficial wounds
- R10** Crushed, degloved, mangled, or pulseless injured extremity
- R11** Two or more proximal long bone fracture sites

- R1** GCS  $\leq 13$  due to trauma
- R2** Active airway assistance required (ie. more than supplemental O2)
- R3** No radial pulse AND heart rate  $\geq 120$
- R4** BP <90 systolic
- R5** Pelvic instability or Chest wall instability or crepitus
- R6** Acute paralysis, loss of sensation, or suspected spinal cord injury
- R7** Amputation proximal to wrist or ankle
- R8**  $\geq 10\%$  BSA partial/full thickness burns
- R9** Penetrating injury to head (or depressed skull fracture), neck, torso, extremities proximal to elbow or knee, excluding superficial wounds
- R10** Crushed, degloved, mangled, or pulseless injured extremity
- R11** Two or more proximal long bone fracture sites

- R1** GCS  $\leq 13$  or change in baseline due to trauma
- R2** Active airway assistance required (ie. more than supplemental O2)
- R3** No radial pulse
- R4** BP <110 systolic
- R5** Pelvic instability or Chest wall instability or crepitus
- R6** Acute paralysis, loss of sensation, or suspected spinal cord injury
- R7** Amputation proximal to wrist or ankle
- R8**  $\geq 5\%$  BSA partial/full thickness burns
- R9** Penetrating injury to head (or depressed skull fracture), neck, torso, extremities proximal to elbow or knee, excluding superficial wounds
- R10** Crushed, degloved, mangled, or pulseless injured extremity
- R11** Two or more proximal long bone fracture sites

Pedi  $\leq 17$

Adult  $\geq 18, < 65$

Geri  $\geq 65$

**If one Blue Criteria met, transport to L3 or L4 Trauma Center;  
OR, if two or more Blue Criteria met, transport to L1 or L3 Trauma Center.**

- B1** Reliable history of any LOC and/or amnesia
- B2**
- B3**
- B4**
- B5** Pregnancy >20 weeks
- B6** Single closed long bone fracture site
- B7** Falls >2X child's height or >10 feet
- B8**
- B9** Ejection from vehicle (excludes open vehicles)
- B10** Driver w/deformed steering wheel
- B11** Death in the same vehicle
- B12** Pedestrian or bicyclist struck; or motorcyclist thrown, run over, or w/significant impact
- B13**
- B14** Weight <10Kg (<22lbs) or RED or PURPLE Broselow Tape Zone
- B15** Suspicion of non-accidental trauma

- B1** Reliable loss of consciousness >5 min.
- B2** Sustained respiratory rate  $\geq 30$  or  $\leq 10$
- B3** Sustained heart rate  $\geq 120$  (w/radial pulse) and BP  $\geq 90$  systolic
- B4** Best motor response = 5
- B5** Pregnancy >20 weeks
- B6** Fracture to humerus or femur due to motor vehicle crash
- B7** Fall from  $\geq 20$  feet
- B8**
- B9** Ejection from vehicle (excludes open vehicles)
- B10** Driver w/deformed steering wheel
- B11** Death in same vehicle
- B12** Pedestrian or bicyclist struck; or motorcyclist thrown, run over, or w/significant impact
- B13** Patient on anticoagulant w/suspected TBI\*
- B14**
- B15**

- B1** Reliable loss of consciousness >5 min.
- B2** Sustained respiratory rate  $\geq 30$  or  $\leq 10$
- B3** Sustained heart rate  $\geq 100$
- B4** Best motor response = 5
- B5**
- B6** Fracture to humerus or femur due to motor vehicle crash
- B7** Fall from  $\geq 3$  feet
- B8** Age  $\geq 65$
- B9** Ejection from vehicle (excludes open vehicles)
- B10** Driver w/deformed steering wheel
- B11** Death in same vehicle
- B12** Pedestrian or bicyclist struck; or motorcyclist thrown, run over, or w/significant impact
- B13** Patient on anticoagulant w/suspected TBI\* (includes daily ASA use)
- B14**
- B15** Significant injuries to two or more body-systems

**\*\*Paramedic intuition may serve as Red/Blue Criteria override.**

**\*Signs/Symptoms of TBI include:**

- Witnessed or reported LOC
- Dizziness, vertigo, or 'lightheadedness'
- Nausea or vomiting
- Changes in vision, photophobia or double vision
- Ataxia or new problems walking, standing, or maintaining balance
- Change in mental status, level of functioning or speech quality